

# Specialist Help

If you feel that more professional help is needed, the general practitioner should be able to tell you what help is available locally, and make a referral to your local child and adolescent mental health service.

The aim is to discover the causes of the problems. It is usual for parents or carers to be involved in treatment. This makes it easier to understand the background to what has happened and to work out what sort of help is needed.

There are different talking approaches, depending on what is causing the problem. Treatment often involves both individual and family work. Individuals will need help with how to cope with the very difficult feelings that cause self-harm. Families often need help in working out how to make sure that the dangerous behavior doesn't happen again, and how to give the support that is needed. If depression or anxiety is part of the problem, these will need treating.

Occasionally, intensive help may be needed. Sometimes, recovery from very damaging or traumatic experiences happens slowly. Then specialist help is needed over a longer period of time.



## USEFUL CONTACTS / HELPFUL ORGANIZATIONS

**SAMARITANS** - 24 hour confidential phone line for people who are in despair and need someone to talk to.  
Tel: 1850 60 90 90 email: Jo@samaritans.org

**AWARE (DEPRESSION AWARENESS)** - operates a helpline service, providing a listening ear for people in distress and their families. (10am—10pm)  
Tel: 1890 303 302

**CHILDLINE** - offers 24 hour support for children and young people in distress. Tel: 1800 66 66 66

**BODYWHYS** - Bodywhys offers support, information and understanding for people with eating disorders, their families and friends  
Tel: 1890 200 444 email: alex@bodywhys.ie

*This leaflet was adapted from information from the Royal College of Psychiatrists.*

*Developed by the  
Regional Suicide Resource Office  
of the  
Health Service Executive - South*

*Supported by the  
National Office for Suicide Prevention*

Ref No. 10-06-0051

Print & Design by Modern Printers Kilkenny

# DELIBERATE SELF HARM IN YOUNG PEOPLE



**Information for  
Parents/Guardian's,  
Teachers etc**

**HE** Feidhmeannacht na Seirbhíse Sláinte  
Health Service Executive

# What is deliberate self harm?

Deliberate self harm is a term used when someone injures or harms themselves on purpose. Common examples include “over-dosing” (self poisoning), hitting, cutting or burning oneself, pulling hair or picking skin, or self strangulation. It can also include taking illegal drugs and excessive amounts of alcohol. Self-harm is always a sign of something being seriously wrong.

# Why do young people harm themselves?

Self-injury is a way of dealing with very difficult feelings that build up inside. People say different things about why they do it;

- Some say that they have been feeling desperate about a problem and don't know where to turn for help. They feel trapped and helpless. Self injury helps them to feel more in control.
- Some people talk of feelings of anger or tension that get bottled up inside, until they feel like exploding. Self injury helps to relieve the tension that they feel
- Feelings of guilt or shame may also become unbearable. Self-harm is a way of punishing oneself.
- Some people try to cope with very upsetting experiences such as trauma or abuse, by convincing themselves that the upsetting event(s) never happened. These people sometimes suffer from feelings of “numbness” or “deadness”.

They say that they feel detached from the world and their bodies, and that self injury is a way of feeling more connected and alive



## Who is at risk?

Self-harm is most commonly triggered by an argument with a parent or close friend. When the young person has experienced abuse, neglect or rejection, they are more likely to harm themselves. Young people who are depressed, or have an eating disorder, are at greater risk. So are people who take illegal drugs or excessive amounts of alcohol.

## Where can I get help?

Anyone who is harming themselves is struggling to cope and needs help. Self injury is often kept secret—even from friends or family. The person feels so ashamed, guilty or bad that they can't face talking about it. There may be clues, such as refusing to wear short sleeves or to take off clothing for sports.

### If you are a parent or teacher, you can help by;

- Recognizing signs of distress, and finding some way of talking with the young person about how they are feeling.
- Listening to their worries and problems, and taking them seriously.
- Offering sympathy and understanding
- Helping with solving problems

- Being clear about the risks of self harm making sure they know that, with help, it will be possible to stop once the underlying problems have been sorted out.
- Making sure that they get the right kind of help as soon as possible.

Its important to make sure that the young person feels that they have someone they can talk to and get support from, when they need it. Otherwise, there is a risk they will harm themselves instead. It's important to ask whether parents and family will be able to give the support that's needed. This may be difficult if there are a lot of problems or arguments at home. As a parent, you may be too upset or angry to be able to give the help that is needed. If so, you should seek advice from your general practitioner.

If you are a teacher, it is important to encourage students to let you know if one of their group is in trouble, upset or shows signs of harming themselves. Friends often worry about betraying a confidence and you may need to explain that self harm can endanger their lives. For this reason, it should never be kept secret.

